

References for Multiple Sclerosis (MS) article from www.drpawluk.com

- Bandari DS, Sternaman D, Chan T, Prostko CR, Sapir T. Evaluating risks, costs, and benefits of new and emerging therapies to optimize outcomes in multiple sclerosis. *J Manag Care Pharm*. 2012 Nov-Dec;18(9):1-17.
- Broła W, Wegrzyn W, Czernicki J. Effect of variable magnetic field on motor impairment and quality of life in patients with multiple sclerosis. *Wiad Lek*. 2002;55(3-4):136-43.
- Capone F, Dileone M, Profice P, Pilato F, Musumeci G, Minicuci G, Ranieri F, Cadossi R, Setti S, Tonali PA, Di Lazzaro V. Does exposure to extremely low frequency magnetic fields produce functional changes in human brain? *J Neural Transm*. 2009 Mar;116(3):257-65.
- Centonze D, Koch G, Versace V, Mori F, Rossi S, Brusa L, Grossi K, Torelli F, Prosperetti C, Cervellino A, Marfia GA, Stanzione P, Marciani MG, Boffa L, Bernardi G. Repetitive transcranial magnetic stimulation of the motor cortex ameliorates spasticity in multiple sclerosis. *Neurology*. 2007 Mar 27;68(13):1045-50.
- Centonze D, Petta F, Versace V, Rossi S, Torelli F, Prosperetti C, Rossi S, Marfia GA, Bernardi G, Koch G, Miano R, Boffa L, Finazzi-Agrò E. Effects of motor cortex rTMS on lower urinary tract dysfunction in multiple sclerosis. *Mult Scler*. 2007 Mar;13(2):269-71.
- de Carvalho ML, Motta R, Konrad G, Battaglia MA, Brichetto G. A randomized placebo-controlled cross-over study using a low frequency magnetic field in the treatment of fatigue in multiple sclerosis. *Mult Scler*. 2012 Jan;18(1):82-9.
- Guseo A. Pulsing electromagnetic field therapy of multiple sclerosis by the Gyuling–Bordacs device: Double-blind, cross-over and open studies. *J Bioelectr* 1987; 6: 23–25.
- Heusser K, Telschaft D and Thoss F. Influence of an alternating 3 Hz magnetic field with an induction of 0.1 millitesla on chosen parameters of the human occipital EEG. *Neurosci Lett* 1997; 239: 57–60.
- Koch G, Rossi S, Prosperetti C, Codecà C, Monteleone F, Petrosini L, Bernardi G, Centonze D. Improvement of hand dexterity following motor cortex rTMS in multiple sclerosis patients with cerebellar impairment. *Mult Scler*. 2008 Aug;14(7):995-8.
- Lappin MS, Lawrie FW, Richards TL and Kramer ED. Effects of a pulsed electromagnetic therapy on multiple sclerosis fatigue and quality of life: A double-blind, placebo controlled trial. *Altern Ther Health Med* 2003; 9: 38–48.
- Mills KR. Magnetic brain stimulation: a tool to explore the action of the motor cortex on single human spinal motoneurons. *Trends Neurosci* 1991 Sep;14(9):401-5.
- Mix E, Henssen HL, Lehmitz R, Lakner K, Hitzschke B, Richter M, Heydenreich A, Meyer-Rienecker H. Pulsing electromagnetic field (PEMF) treatment of patients with multiple sclerosis (MS) and migraine. Effect on cell volume and phagocytic activity of blood cells and clinical results. *J Bioelectr* 1990; 9(2):242. Third Symposium on Magnetotherapy and Magnetic Stimulation, October 1989, Szekesfehervar, Hungary.
- Mostert S and Kesselring J. Effect of pulsed magnetic field therapy on the level of fatigue in patients with multiple sclerosis – a randomized controlled trial. *Mult Scler* 2005; 11: 302–305.
- Nielsen JF, Sinkjaer T and Jakobsen J. Treatment of spasticity with repetitive magnetic stimulation; a double-blind placebo-controlled study. *Mult Scler* 1996; 2: 227–232.
- Piatkowski J, Kern S and Ziemssen T. Effect of BEMER magnetic field therapy on the level of fatigue in patients with multiple sclerosis: a randomized double – blind controlled trial. *J Altern Complement Med* 2009; 15: 507–511.
- Quittan M, Schuhfried O, Wiesinger GF and Fialka-Moser V. Clinical effectiveness of magnetic field therapy a review of the literature. *Acta Med Austriaca* 2000; 27: 61–68.

- Ravnborg M, Knudsen GM, Blinkenberg M. No effect of pulsed magnetic stimulation on the blood-brain barrier in rats. *Neuroscience* 38(1):277-280, 1990.
- Richards TL, Lappin MS, Acosta-Urquidi J, et al. Double-blind study of pulsing magnetic field effects on multiple sclerosis. *J Altern Complement Med* 1997; 3: 21–29.
- Sandyk R. Application of weak electromagnetic fields facilitates sensory–motor integration in patients with multiple sclerosis. *Int J Neurosci* 1996; 85: 101–110.
- Sandyk R. Resolution of sleep paralysis by weak electromagnetic fields in a patient with multiple sclerosis. *Int J Neurosci* 1997 Aug;90(3-4):145-57.
- Sandyk R. Serotonergic neuronal atrophy with synaptic inactivation, not axonal degeneration, are the main hallmarks of multiple sclerosis. *Int J Neurosci* 1998 Jul;95(1-2):133-40.
- Sandyk R. Treatment with electromagnetic fields reverses the long-term clinical course of a patient with chronic progressive multiple sclerosis. *Int J Neurosci* 1997 Aug;90(3-4):177-85.
- Sandyk R. Treatment with weak electromagnetic fields improves fatigue associated with multiple sclerosis. *Int J Neurosci* 1996; 84: 177–186.
- Sandyk R. Weak electromagnetic fields potentiate the effects of 4-aminopyridine in multiple sclerosis. *Int J Neurosci* 1996; 85: 125–129.
- Sandyk R. Yawning and stretching--a behavioral syndrome associated with transcranial application of electromagnetic fields in multiple sclerosis. *Int J Neurosci* 1998 Jul;95(1-2):107-13.
- Terlaki, G.; Gyorgy, B.; Payer, A. Clinical experiences by magnetotherapy. *Hungarian Symposium on Magnetotherapy (with International Participation), 2nd symposium, 16-17 May, Szekesfehervar, Hungary, p. 175-179, 1987.*
- Wieczorek V, Schock G. Our experience with the pulsing electromagnetic field therapy of neurologic diseases. *J Bioelectr* 1990; 9(2):242 Third Symposium on Magnetotherapy and Magnetic Stimulation, October 1989, Szekesfehervar, Hungary.
- Yamamoto D, Campbell JD. Cost-effectiveness of multiple sclerosis disease-modifying therapies: a systematic review of the literature. *Autoimmune Dis.* 2012;2012:784364.