

Urinary Incontinence and OAB

By Dr. W. Pawluk

References

- Ammi M, Chautard D, Brassart E, Culty T, Azzouzi AR, Bigot P. Transcutaneous posterior tibial nerve stimulation: evaluation of a therapeutic option in the management of anticholinergic refractory overactive bladder. *Int Urogynecol J*. 2014 Aug;25(8):1065-9.
- Bartley J, Gilleran J, Peters K. Neuromodulation for overactive bladder. *Nat Rev Urol*. 2013 Sep;10(9):513-21.
- But I, Varda NM. Functional magnetic stimulation: a new method for the treatment of girls with primary nocturnal enuresis? *J Pediatr Urol*. 2006 Oct;2(5):415-8.
- But I. Conservative treatment of female urinary incontinence with functional magnetic stimulation. *Urology*. 2003 Mar;61(3):558-61.
- Eftekhari T, Teimoori N, Miri E, Nikfallah A, Naeimi M, Ghajarzadeh M.
- Fujishiro T, Enomoto H, Ugawa Y, Takahashi S, Ueno S, Kitamura T. Magnetic stimulation of the sacral roots for the treatment of stress incontinence: an investigational study and placebo controlled trial. *J Urol*. 2000 Oct;164(4):1277-9.
- Fujishiro T, Takahashi S, Enomoto H, Ugawa Y, Ueno S, Kitamura T. Magnetic stimulation of the sacral roots for the treatment of urinary frequency and urge incontinence: an investigational study and placebo controlled trial. *J Urol*. 2002 Sep;168(3):1036-9.
- Gaspard L, Tombal B, Opsomer RJ, Castille Y, Van Pesch V, Detrembleur C. Physiotherapy and neurogenic lower urinary tract dysfunction in multiple sclerosis patients: a randomized controlled trial. *Prog Urol*. 2014 Sep;24(11):697-707.
- Gaziev G, Topazio L, Iacovelli V, Asimakopoulos A, Di Santo A, De Nunzio C, Finazzi-Agrò E. Percutaneous Tibial Nerve Stimulation (PTNS) efficacy in the treatment of lower urinary tract dysfunctions: a systematic review. *BMC Urol*. 2013 Nov 25;13:61.
- Gungor Ugurlucan F, Onal M, Aslan E, Ayyildiz Erkan H, Kizilkaya Beji N, Yalcin O. Comparison of the effects of electrical stimulation and posterior tibial nerve stimulation in the treatment of overactive bladder syndrome. *Gynecol Obstet Invest*. 2013;75(1):46-52.
- Hartmann KE, McPheeters ML, Biller DH, Ward RM, McKoy JN, Jerome RN, Micucci SR, Meints L, Fisher JA, Scott TA, Slaughter JC, Blume JD. Treatment of overactive bladder in women. *Evid Rep Technol Assess (Full Rep)*. 2009 Aug;(187):1-120.

- Ishikawa N, Suda S, Sasaki T, Yamanishi T, Hosaka H, Yasuda K, Ito H. Development of a non-invasive treatment system for urinary incontinence using a functional continuous magnetic stimulator (FCMS). *Med Biol Eng Comput.* 1998 Nov;36(6):704-10.
- Janssen DA, Farag F, Heesakkers JP. Urgent-SQ implant in treatment of overactive bladder syndrome: 9-year follow-up study. *Neurourol Urodyn.* 2013 Jun;32(5):472-5.
- Manganotti P, Zaina F, Vedovi E, Pistoia L, Rubilotta E, D'Amico A, Falso M. Eura Repetitive magnetic stimulation of the sacral roots for the treatment of stress incontinence: a brief report. *Medicophys.* 2007 Sep;43(3):339-44.
- Martinson M, MacDiarmid S, Black E. Cost of neuromodulation therapies for overactive bladder: percutaneous tibial nerve stimulation versus sacral nerve stimulation. *J Urol.* 2013 Jan;189(1):210-6.
- Monteiro ÉS, de Carvalho LB, Fukujima MM, Lora MI, do Prado GF. Electrical stimulation of the posterior tibialis nerve improves symptoms of poststroke neurogenic overactive bladder in men: a randomized controlled trial. *Urology.* 2014 Sep;84(3):509-14.
- Peters KM, Carrico DJ, Wooldridge LS, Miller CJ, MacDiarmid SA. Percutaneous tibial nerve stimulation for the long-term treatment of overactive bladder: 3-year results of the STEP study. *J Urol.* 2013 Jun;189(6):2194-201.
- Posterior tibial nerve stimulation for treating neurologic bladder in women: a randomized clinical trial. *Acta Med Iran.* 2014;52(11):816-21.
- Sato E, Ueda Y, Imai Y, Suda S, Nakamura T, Yamanishi T, Shinoda M. Pulsed magnetic stimulation with a high-frequency continuous magnetic stimulator (SMN-X) does not exert an adverse effect on genital organs and the estrous cycle in female lar:Wistar-Imamichi rats. *Neurourol Urodyn.* 2011 Nov;30(8):1675-80.
- Shalom DF, Pillalamarri N, Xue X, Kohn N, Lind LR, Winkler HA, Metz CN. Sacral nerve stimulation reduces elevated urinary nerve growth factor levels in women with symptomatic detrusor overactivity. *Am J Obstet Gynecol.* 2014 Nov;211(5):561.e1-5.
- Sun HD, Chung SD. Could overactive bladder be a chronic inflammatory disorder? *Incont Pelvic Floor Dysfunct* 2009; 3(Suppl 1):17-19.
- Suzuki T, Yasuda K, Yamanishi T, Kitahara S, Nakai H, Suda S, Ohkawa H. Randomized, double-blind, sham-controlled evaluation of the effect of functional continuous magnetic stimulation in patients with urgency incontinence. *Neurourol Urodyn.* 2007;26(6):767-72.
- Wallis MC, Davies EA, Thalib L, Griffiths S. Pelvic static magnetic stimulation to control urinary incontinence in older women: A randomized controlled trial. *Clin Med Res.* 2011 Aug 4.

- Yamanishi T, Homma Y, Nishizawa O, Yasuda K, Yokoyama O. Multicenter, randomized, sham-controlled study on the efficacy of magnetic stimulation for women with urgency urinary incontinence. *Int J Urol*. 2014 Apr;21(4):395-400.
- Yamanishi T, Sakakibara R, Uchiyama T, Suda S, Hattori T, Ito H, Yasuda K. Comparative study of the effects of magnetic versus electrical stimulation on inhibition of detrusor overactivity. *Urology*. 2000 Nov 1;56(5):777-81.
- Yamanishi T, Yasuda K, Sakakibara R, Suda S, Ishikawa N, Hattori T, Hosaka H. Induction of urethral closure and inhibition of bladder contraction by continuous magnetic stimulation. *Neurourol Urodyn*. 1999;18(5):505-10.
- Yamanishi T, Yasuda K, Suda S, Ishikawa N, Sakakibara R, Hattori T. Effect of functional continuous magnetic stimulation for urinary incontinence. *J Urol*. 2000 Feb;163(2):456-9.
- Yamanishi T1, Kamai T, Yoshida K. Neuromodulation for the treatment of urinary incontinence. *Int J Urol*. 2008 Aug;15(8):665-72.