

References for Concussion/TBI Article on www.drpawluk.com

- Anderson B, Mishory A, Nahas Z, Borckardt JJ, Yamanaka K, Rastogi K, George MS. Tolerability and safety of high daily doses of repetitive transcranial magnetic stimulation in healthy young men. *J ECT*. 2006 Mar;22(1):49-53.
- Baker-Price L, Persinger MA. Intermittent burst-firing weak (1 microTesla) magnetic fields reduce psychometric depression in patients who sustained closed head injuries: a replication and electroencephalographic validation. *Percept Mot Skills*. 2003 Jun;96(3 Pt 1):965-74.
- Baker-Price LA, Persinger MA. Weak, but complex pulsed magnetic fields may reduce depression following traumatic brain injury. *Percept Mot Skills*. 1996 Oct;83(2):491-8.
- Bartko D, Turcáni P, Danisová J, et al. The effects of the pulsing magnetic field on the cerebral circulation, eeg power spectra and some properties of the blood. A preliminary data. *J Bioelectr* 7(1):131-132, 1988.
- Bartnik-Olson BL, Harris NG, Shijo K, Sutton RL. Insights into the metabolic response to traumatic brain injury as revealed by (13)C NMR spectroscopy. *Front Neuroenergetics*. 2013 Oct 4;5:8.
- Başar E, Schürmann M, Başar-Eroglu C, Karakaş S. Alpha oscillations in brain functioning: an integrative theory. *Int J Psychophysiol*. 1997 Jun;26(1-3):5-29.
- Bell, G., A. Marino, A. Chesson and F. Struve (1991) Human sensitivity to weak magnetic fields. *Lancet*, 338: 1521—1522.
- Berrigan L, Marshall S, McCullagh S, et al. Quality of clinical practice guidelines for persons who have sustained mild traumatic brain injury. *Brain Inj*. 2011;25(7-8):742-51.
- Bharath RD, Munivenkatappa A, Gohel S, et al. Recovery of resting brain connectivity ensuing mild traumatic brain injury. *Front Hum Neurosci*. 2015 Sep 22;9:513.
- Blackman CF. Stimulation of brain tissue in vitro by extremely low frequency, low intensity, sinusoidal electromagnetic fields. *Prog Clin Biol Res* 257:107-117. *Electromagnetic Fields and Neurobehavioral Function*, M. E. O'Connor and R. H. Lovely, eds., New York: Alan R. Liss, Inc. 1988.
- Bonni S, Mastropasqua C, Bozzali M, et al. Theta burst stimulation improves visuo-spatial attention in a patient with traumatic brain injury. *Neurol Sci*. 2013 Nov;34(11):2053-6.
- Choe MC, Giza CC. Diagnosis and management of acute concussion. *Semin Neurol*. 2015 Feb;35(1):29-41.
- Clark VP, Parasuraman R. Neuroenhancement: enhancing brain and mind in health and in disease. *Neuroimage*. 2014 Jan 15;85 Pt 3:889-94.
- Cosentino G, Giglia G, Palermo A, et al. A case of post-traumatic complex auditory hallucinosis treated with rTMS. *Neurocase*. 2010 Jun;16(3):267-72.

- Del Percio C, Marzano N, Tilgher S, Fiore A, Di Ciolo E, Aschieri P, Lino A, Toràn G, Babiloni C, Eusebi F. Pre-stimulus alpha rhythms are correlated with post-stimulus sensorimotor performance in athletes and non-athletes: a high-resolution EEG study. *Clin Neurophysiol.* 2007 Aug;118(8):1711-20.
- Diamond MC, Tenforde TS, Liburdy RP, et al. The influence of ultrahigh magnetic fields on cerebral cortical morphological development: a preliminary study (meeting abstract). *Bioelectromagnetics Society, 11th Annual Meeting, 18-22 June, Tucson, AZ, Abstract No. P-1-20, p. 62-63, 1989.*
- Dowman R, Wolpaw JR, Seegal RF, Satya-Murti S. Chronic exposure of primates to 60-Hz electric and magnetic fields: III. Neurophysiologic effects. *Bioelectromagnetics.* 1989;10(3):303-17.
- Esty ML and Nelson D. Neurotherapy of TBI/PTSD in OEF/OIF Veterans. *The Journal of Neuropsychiatry and Clinical Neurosciences,* 21:221-223, 2009.
- Gavalas RJ, Walter DO, Hamer J, Adey WR. Effect of low-level, low-frequency electric fields on EEG and behavior in *Macaca nemestrina.* *Brain Res* 1970; 18 (3): 491 – 501.
- George MS, Raman R, Benedek DM, et al. A two-site pilot randomized 3 day trial of high dose left prefrontal repetitive transcranial magnetic stimulation (rTMS) for suicidal inpatients. *Brain Stimul.* 2014 May-Jun;7(3):421-31.
- Goodwin T. Physiological and molecular genetic effects of time-varying electromagnetic fields on human neuronal cells. NASA Johnson Space Center, Houston, TX, United States. NASA/TP-2003-212054.
- Grunner O. Cerebral use of a pulsating magnetic field in neuropsychiatry patients with long-term headache. *EEG EMG Z Elektroenzephalogr Verwandte Geb* (1985) Dec;16(4):227-230.
- Herrold AA, Kletzel SL, Harton BC, et al. Transcranial magnetic stimulation: potential treatment for co-occurring alcohol, traumatic brain injury and posttraumatic stress disorders. *Neural Regen Res.* 2014 Oct 1;9(19):1712-30.
- Hoffer ME. Mild traumatic brain injury: neurosensory effects. *Curr Opin Neurol.* 2015 Feb;28(1):74-7.
- Ingram DA, Thompson AJ, Swash M. Central motor conduction in multiple sclerosis: evaluation of abnormalities revealed by transcutaneous magnetic stimulation of the brain. *J Neurol Neurosurg Psychiatry* 51(4):487-494, 1988.
- Jin Y, Phillips B. A pilot study of the use of EEG-based synchronized Transcranial Magnetic Stimulation (sTMS) for treatment of Major Depression. *BMC Psychiatry.* 2014 Jan 18;14:13.
- Koski L, Kolivakis T, Yu C, et al. Noninvasive brain stimulation for persistent postconcussion symptoms in mild traumatic brain injury. *J Neurotrauma.* 2015 Jan 1;32(1):38-44.

- Lee J, Choi BH, Oh E, et al. Treatment of Alzheimer's Disease with Repetitive Transcranial Magnetic Stimulation Combined with Cognitive Training: A Prospective, Randomized, Double-Blind, Placebo-Controlled Study. *J Clin Neurol*. 2015 Sep 11.
- Leuchter AF, Cook IA, Feifel D, et al. Efficacy and Safety of Low-field Synchronized Transcranial Magnetic Stimulation (sTMS) for Treatment of Major Depression. *Brain Stimul*. 2015;8(4):787-94.
- Leuchter AF, Cook IA, Jin Y, Phillips B. The relationship between brain oscillatory activity and therapeutic effectiveness of transcranial magnetic stimulation in the treatment of major depressive disorder. *Front Hum Neurosci*. 2013 Feb 26;7:37.
- Leuchter AF, Hunter AM, Krantz DE, Cook IA. Rhythms and blues: modulation of oscillatory synchrony and the mechanism of action of antidepressant treatments. *Ann N Y Acad Sci*. 2015 May;1344:78-91.
- Leung A, Fallah A, Shukla S, Lin L, Tsia A, Song D, Polston G, Lee R. rTMS in Alleviating Mild TBI Related Headaches - A Case Series. *Pain Physician*. 2016 Feb;19(2):E347-54.
- Liao X, Li G, Wang A, et al. Repetitive Transcranial Magnetic Stimulation as an Alternative Therapy for Cognitive Impairment in Alzheimer's Disease: A Meta-Analysis. *J Alzheimers Dis*. 2015 Sep 9;48(2):463-72.
- Ling H, Hardy J, Zetterberg H. Neurological consequences of traumatic brain injuries in sports. *Mol Cell Neurosci*. 2015 May;66(Pt B):114-22.
- Louise-Bender Pape T, Rosenow J, Lewis G, et al. Repetitive transcranial magnetic stimulation-associated neurobehavioral gains during coma recovery. *Brain Stimul*. 2009 Jan;2(1):22-35.
- Lustenberger C, Boyle MR, Foulser AA, Mellin JM, Fröhlich F. Functional role of frontal alpha oscillations in creativity. *Cortex*. (2015) Jun;67:74-82.
- MacFarlane MP, Glenn TC. Neurochemical cascade of concussion. *Brain Inj*. 2015;29(2):139-53.
- Marshall S, Bayley M, McCullagh S, et al. Clinical practice guidelines for mild traumatic brain injury and persistent symptoms. *Can Fam Physician*. 2012 Mar;58(3):257-67, e128-40.
- Meehan WP 3rd. Medical therapies for concussion. *Clin Sports Med*. 2011 Jan;30(1):115-24, ix.
- Mez J, Solomon TM, Daneshvar DH, et al. Pathologically Confirmed Chronic Traumatic Encephalopathy in a 25-Year-Old Former College Football Player. *JAMA Neurol*. 2016 Jan 4:1-3.
- Nelson DV, and Esty ML. Neurotherapy for pain in veterans with trauma spectrum disorders. *The Journal of Pain* 10:S18, 2009.
- Nelson DV, Esty ML. Neurotherapy for chronic headache following traumatic brain injury. *Mil Med Res*. 2015 Aug 31;2:22.

- Nelson, D and Esty, ML. Neurotherapy for Chronic TBI/PTSD Symptoms in Vietnam Veterans. *The Journal of Head Trauma Rehabilitation*. 2009 (24)5, 403.
- Niedermeyer, E. (1999) *The Normal EEG of the Waking Adult*. Electroencephalography. Lippincott Williams y Wilkins, Baltimore.
- Nielson DM, McKnight CA, Patel RN, et al. Preliminary guidelines for safe and effective use of repetitive transcranial magnetic stimulation in moderate to severe traumatic brain injury. *Arch Phys Med Rehabil*. 2015 Apr;96(4 Suppl):S138-44.
- Ontario Neurotrauma Foundation. *Guidelines for concussion/mild traumatic brain injury and persistent symptoms*. 2nd edition. 2013.
- O'Reardon JP, Solvason HB, Janicak PG, et al. Efficacy and safety of transcranial magnetic stimulation in the acute treatment of major depression: a multisite randomized controlled trial. *Biol Psychiatry*. 2007 Dec 1;62(11):1208-16.
- Ossenkopp KP, Cain DP. Inhibitory effects of acute exposure to low-intensity 60-hz magnetic fields on electrically kindled seizures in rats. *Brain Res* 442(2):255-260, 1988.
- Ossenkopp KP, Kavaliers M. Clinical and applied aspects of magnetic field exposure: a possible role for the endogenous opioid system. *J Bioelectr* 7(2):189-208, 1989.
- Pape TL, Rosenow J, Lewis G. Transcranial magnetic stimulation: a possible treatment for TBI. *J Head Trauma Rehabil*. 2006 Sep-Oct;21(5):437-51.
- Persinger MA, Saroka KS. Comparable proportions of classes of experiences and intracerebral consequences for surgical stimulation and external application of weak magnetic field patterns: implications for converging effects in complex partial seizures. *Epilepsy Behav*. 2013 Apr;27(1):220-4.
- Peskind ER, Brody D, Cernak I, et al. Military- and sports-related mild traumatic brain injury: clinical presentation, management, and long-term consequences. *J Clin Psychiatry* (2013)74: 180–188.
- Politis MJ, Zanakis MF. Treatment of the damaged rat hippocampus with a locally applied electric field. *Exp Brain Res* 71(1):223-226, 1988.
- Raji CA, Tarzwell R, Pavel D, et al. Clinical utility of SPECT neuroimaging in the diagnosis and treatment of traumatic brain injury: a systematic review. *PLoS One*. 2014 Mar 19;9(3):e91088.
- Rasouli J, Lekhraj R, White NM, Flamm ES, Pilla AA, Strauch B, Casper D. Attenuation of interleukin-1beta by pulsed electromagnetic fields after traumatic brain injury. *Neurosci Lett*. 2012 Jun 21;519(1):4-8.
- Reti IM, Schwarz N, Bower A, et al. Transcranial magnetic stimulation: A potential new treatment for depression associated with traumatic brain injury. *Brain Inj*. 2015;29(7-8):789-97.

- Rohan ML, Yamamoto RT, Ravichandran CT, et al. Rapid mood-elevating effects of low field magnetic stimulation in depression. *Biol Psychiatry*. 2014 Aug 1;76(3):186-93.
- Rossia S, Hallett M, Rossini, PM, Pascual-Leone A. The Safety of TMS Consensus Group¹. Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clin Neurophysiol*. 2009 December ; 120(12): 2008–2039.
- Salam MT, Kassiri H, Genov R, et al. Rapid brief feedback intracerebral stimulation based on real-time desynchronization detection preceding seizures stops the generation of convulsive paroxysms. *Epilepsia*. 2015 Aug;56(8):1227-38.
- Salford LG, Brun A, Eberhardt JL, Persson BRR. Development of rat brain tumours during exposure to continuous and pulsed 915 MHz electromagnetic radiation (meeting abstract). First World Congress for Electricity and Magnetism in Biology and Medicine, 14-19 June, Lake Buena Vista, FL, Abstract No. I-1, p. 27-28, 1992.
- Schoenberger NE, Shif SC, Esty ML, et al. Flexyx Neurotherapy System in the treatment of traumatic brain injury: an initial evaluation. *J Head Trauma Rehabil*. 2001 Jun;16(3):260-74.
- Seegal RF, Wolpaw JR, Dowman R. Chronic exposure of primates to 60-Hz electric and magnetic fields: II. Neurochemical effects. *Bioelectromagnetics*. 1989;10(3):289-301.
- Sieron A, Labus L, Nowak P, Cieslar G, Brus H, Durczok A, Zagzil T, Kostrzewa RM, Brus R. Alternating extremely low frequency magnetic field increases turnover of dopamine and serotonin in rat frontal cortex. *Bioelectromagnetics* 25:426– 430, 2004.
- Simpson G, Tate R. Suicidality in people surviving a traumatic brain injury: prevalence, risk factors and implications for clinical management. *Brain Inj*. 2007 Dec;21(13-14):1335-51.
- Sokhadze EM, El-Baz AS, Tasman A, et al. Neuromodulation integrating rTMS and neurofeedback for the treatment of autism spectrum disorder: an exploratory study. *Appl Psychophysiol Biofeedback*. 2014 Dec;39(3-4):237-57.
- Tang HY, Vitiello MV, Perlis M, Riegel B. Open-loop neurofeedback audiovisual stimulation: a pilot study of its potential for sleep induction in older adults. *Appl Psychophysiol Biofeedback*. 2015 Sep;40(3):183-8.
- VA/DoD Clinical Practice Guideline for Management of Concussion/Mild Traumatic Brain Injury. Management of Concussion/mTBI Working Group. *J Rehabil Res Dev*. 2009;46(6):CP1-68.
- Warden DL, Bleiberg J, Cameron KL, et al. Persistent prolongation of simple reaction time in sports concussion. *Neurology*. 2001 Aug 14;57(3):524-6.
- Wever, R.A. (1987) The electromagnetic environment and the circadian rhythms of human subjects. In: M. Grandolfo, SM. Michaelson and A. Rindi (Eds.), *Biological Effects and Dosimetry of Static and ELF Electromagnetic Fields*, Plenum Press, New York, NY.

- Wolpaw JR, Seegal RF, Dowman R. Chronic exposure of primates to 60-Hz electric and magnetic fields: I. Exposure system and measurements of general health and performance. *Bioelectromagnetics*. 1989;10(3):277-88.
- Yuh EL, Mukherjee P, Lingsma HF, et al. (2013) Magnetic resonance imaging improves 3-month outcome prediction in mild traumatic brain injury. *Ann Neurol* 73: 224–235.
- Zecca,L, Margonato V, Esposti G, et al. Brain transmitters in rats exposed to 50 hz pulsed magnetic fields. (meeting abstract) *J Bioelectr* 8(2):269, 1989. International Symposium in Honor of Luigi Galvani, 14-16, April, Bologna, Italy.